



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short communication

Comparison of suicide attempts among nationally representative samples of Mexican adolescents 12 months before and after the outbreak of the Covid-19 pandemic

Rosario Valdez-Santiago^a, Aremis Villalobos^{b,*}, Luz Arenas-Monreal^a, Catalina González-Forteza^c, Alicia Edith Hermosillo-de-la-Torre^d, Corina Benjet^c, Fernando A. Wagner^e

^a Center for Health Systems Research, National Institute of Public Health (INSP), Mexico

^b Center for Population Health Research, National Institute of Public Health (INSP), Mexico

^c Department of Epidemiology and Psychosocial Research, National Institute of Psychiatry (INP), Mexico

^d Psychology Department, Autonomous University of Aguascalientes (UAA), Mexico

^e School of Social Work, University of Maryland, Baltimore, USA

ARTICLE INFO

Keywords:

Suicide attempts
Adolescents, Covid-19

ABSTRACT

Background: There is serious concern over the increase in mental health problems during the coronavirus disease 2019 (Covid-19) pandemic.

Methods: Based on data from two Mexican National Health and Nutrition Surveys conducted in 2018–2019 and 2020 ($n = 17,925$ and $4,913$, respectively), we estimated the prevalence of suicide attempts among adolescents 10–19 years old in the previous year. We constructed a multivariate logistic regression model adjusted by sociodemographic characteristics and contextual variables for the Covid-19 pandemic.

Results: The prevalence of suicide attempts in the previous year was similar in both surveys. We found that women, youth in urban localities and individuals living in households where a family member had lost her/his job as a result of the Covid-19 contingency were more likely to attempt suicide compared to their counterparts. On the other hand, attending classes online proved to be a protective factor (aOR=0.3, 95% CI=0.1, 0.8, $p = 0.022$).

Limitations: The principal limitation of our study concerned the restricted size of our sample for the 2020 survey wave.

Conclusions: Population-level policies aimed at providing economic protection and helping youth to return to school would exert a favorable impact on the mental health and suicidal behavior of youths.

1. Introduction

On March 11, 2020, the World Health Organization (WHO) declared Covid-19 a global pandemic (World Health Organization, 2020); 22 days later, Mexico announced a national state of emergency and implemented a series of precautionary measures including home confinement. It is well known that pandemics affect mental health in proportion to the magnitude of the event and levels of population vulnerability. Covid-19 has posed unexpectedly high mental health risks for the world at large (Xiong et al., 2020), and Mexico is no exception. Suicide is a public mental health problem characterized as an unnatural and preventable death.

Suicide attempts (SAs) constitute an important risk factor for death by suicide (Rockstroh et al., 2021). They are multifactorial and have specific characteristics, for instance, women are more likely to attempt suicide than men. In Mexico, SAs have risen for both sexes in the last three national health surveys (Valdez-Santiago et al., 2021), with adolescents and young adults representing a major risk group (Hermosillo-de la Torre et al., 2021). Furthermore, SAs have trended downward with marked differences in the incidence of SAs observed between urban and rural areas (INEGI, 2019).

In light of the characteristics of this public health problem, its rising prevalence, its association with sociodemographic and psychological factors (Hermosillo-de la Torre et al., 2020; Rivera-Rivera et al., 2020)

* Corresponding autor.

E-mail address: alvillalobos@insp.mx (A. Villalobos).

<https://doi.org/10.1016/j.jad.2021.10.111>

Received 3 June 2021; Received in revised form 10 September 2021; Accepted 23 October 2021

Available online 29 October 2021

0165-0327/© 2021 Elsevier B.V. All rights reserved.

and its potential upsurge as a result of confinement, the aim of this report is to compare the prevalence of SAs in the Mexican adolescent population during the 12 months prior and subsequent to the implementation of Covid-19 confinement restrictions. We had no *a priori* hypothesis given the novelty of the pandemic situation. Additionally, our work sought to disaggregate estimates by sex, age and urban/rural area, and to assess the association between SAs and sociodemographic as well as psychosocial factors.

2. Methods

Based on data from the 2018–2019 National Health and Nutrition Survey (*ENSANUT*) as well as the 2020 Covid *ENSANUT* (*ENSANUT-COVID*), we estimated the prevalence of last-year SAs (LYSAs) in adolescents 10–19 years old.

2.1. Sample selection

The 2018–19 *ENSANUT* and 2020 *ENSANUT-COVID* were household surveys with a probabilistic, multi-stage sample design, representative at the national, regional and rural/urban levels ($n = 17,925$ and $4,812$, respectively) (Romero-Martínez et al., 2019; Romero-Martínez et al., 2021). The 2018–19 *ENSANUT* was also representative at the state level; this was impossible to accomplish in 2020 because of the protective measures implemented to prevent Covid-19 infection among field staff and participants, that limited sample size. In both surveys, households were randomly selected, and then an additional list of one randomly selected adolescent per household was compiled. Exclusion criteria consisted of refusing to participate and having auditive or cognitive disabilities that would render individuals unable to respond to the survey. All participants signed informed-consent forms. Our study was approved by the Institutional Review Board of the National Institute of Public Health in Mexico.

2.2. Variables of interest

Data were collected from July 2018 to June 2019 (*ENSANUT 2018,19*) and from August to November 2020 (*ENSANUT-COVID 2020*). Comparable standardized questionnaires for adolescents were administered personally, at the homes, by trained personnel.

2.2.1. Last-year suicide attempts

The term suicide attempt was defined as “any action taken by a given person that could cause them to die.” (Remus, 2020) We created a dichotomous variable for LYSA occurrence based on two questions: (1) “Have you ever attempted to harm yourself or deliberately cut, intoxicated or hurt yourself in any way for the purpose of dying?” and (2) “Was this in the last 12 months?”

2.2.2 Sociodemographic variables included sex, age and type of locality (urban or rural). We generated a socioeconomic indicator grouping households into income deciles; the method used has been described elsewhere (Gutiérrez, 2013).

2.2.3 Covid-19 context indicators included the following: at least one household member had stopped receiving income because of the contingency (yes, no); at least one household member had lost his/her job because of the contingency (yes, no); household members had been forced to reduce their spending during the contingency (yes, no); school-aged household members were attending virtual classes (yes, no).

2.3. Statistical analyses

We tabulated the characteristics of participating adolescents according to whether or not they had attempted suicide, with 95% confidence intervals. Poisson regression was used to estimate the prevalence ratios of SAs. Based on logistic bivariate and multivariate models with a 5% level of significance, we estimate variables associated to last-year

suicide attempts in 10–19-year-old adolescents. The numbers have been rounded to one decimal place. Statistical analyses were performed using STATA SE 14.0 software, with estimates taking into account the probabilistic, multi-stage sample design and unequal selection probabilities.

3. Results

Table 1 depicts LYSA prevalence over 12 months as well as sample characteristics, stratifying the data according to whether the attempt took place prior to or during the survey. Prevalence estimates for were 1.8% (95% Confidence Interval, CI = 1.6, 2.1) prior to Covid-19 and 2.1% during the first month of the pandemic (95% CI = 1.4, 3.1), showing a non-significant statistical difference. Interestingly, the prevalence of SAs decreased among men in 2020 (0.3%, 95% CI = 0.1, 0.8) as compared to 2018–19 (1, 95% CI = 0.7, 1.4), but increased among women (3.8, 95% CI = 2.4, 5.9 in 2020, vs. 2.7, 95% CI = 2.3, 3.2 in 2018–19). Prevalence among youth 10–14 years old was 1.4% in both surveys (95% CI = 1.1, 1.8 and 0.8, 2.6, respectively), but increased slightly among those 15–19 years old in 2020; however, confidence intervals overlapped markedly (2.3% in 2018–19 vs. 2.8% in 2020; 95% CI = 1.9, 2.8 and 1.4, 5.3, respectively). The figures revealed a downward trend in SA prevalence by socioeconomic level: those at the lowest level showed the highest prevalence (2.2% in 2018–19 vs. 2.3% in 2020) and those at the highest level the lowest prevalence (1.6% vs. 1.7%, respectively). However, the corresponding confidence intervals overlapped, indicating a non-significant statistical difference. Finally, while no difference was observed in SA prevalence between youth in rural as opposed to urban areas pre-Covid-19 (1.8%), prevalence in 2020 proved higher among the latter (2.6%, 95% CI = 1.6, 4.1) and lower among the former (0.8%, 95% CI = 0.4, 1.9).

Table 2 illustrates the results of three groups of logistic regression models comparing estimates for the covariates in our analysis. Model 1 included unadjusted estimates. Model 2 was adjusted simultaneously for sociodemographic variables, and Model 3 included fully adjusted estimates for all variables. We found no evidence of major confounding variables, but registered a number of noteworthy associations, namely that women were more likely to attempt suicide than men, with other variables remaining unchanged (adjusted odds ratio, aOR = 12.5, 95% CI = 4.7, 33.4, $p < 0.001$); urban youth were 4.4 times more likely to attempt suicide than their rural counterparts (95% CI = 1.5, 12.9; $p = 0.008$); and SAs were more likely to occur in households where a family member had lost his/her job because of Covid-19 (aOR = 2.8, 95% CI = 1.2, 6.5, $p = 0.014$). However, they were less likely among participants who had been forced to reduce their expenses during the contingency (aOR = 0.5, 95% CI = 0.3, 1.0, $p = 0.039$) as well as in those cases where school-aged –including adolescent– members of the household were attending classes online (aOR = 0.3, 95% CI = 0.1, 0.8, $p = 0.022$).

4. Discussion

We found no significant difference in the prevalence of SAs among adolescents prior to and during the pandemic, with a pre-pandemic rate of 1.8% compared to 2.1% during the pandemic period. This is consistent with a study of 21 mostly high-income countries, which found no evidence of an increase in suicide deaths among the total population during the initial months of the Covid-19 pandemic (Pirkis et al., 2021). Mexico was one of the few upper-middle-income countries included in this study. A meta-analysis of 54 studies found a 2.68% SA rate for all age groups in community samples during the pandemic. While females, younger individuals and those in less democratic countries were more vulnerable to suicidal ideation during the pandemic, only the level of democracy was associated with actual SAs (Dubé et al., 2021).

While there may have been no overall increase in SAs among adolescents in Mexico during this period, some individuals in this age group were at greater risk than others. We observed that female adolescents (as

the topic during the pandemic period in Latin America. However, it is important to note that, because of the 12-month recall period and the dates of fieldwork, our estimation of SAs during the pandemic period covered only the first five to eight months of the pandemic, depending on when respondents were surveyed. Therefore, if SAs truly increased during the pandemic, we would most likely have underestimated SA prevalence; this may account for the lack of statistically significant differences. On the other hand, if SAs truly decreased during the pandemic, we would most likely have overestimated SA prevalence. Additionally, our conclusions were limited by the cross-sectional survey design that yielded no evidence for establishing the causality or directionality of the associations between SAs and their correlates. Finally, measuring SAs depended on the self-reporting of a sensitive issue burdened with cultural and religious taboos. Hence, our prevalence estimates were likely to be conservative.

6. Conclusion

While there may have been no overall increase in SAs among Mexican adolescents during the first five to eight months of the pandemic, adolescents in households that lost employment and those not attending on-line classes are at risk, exacerbating the already large mental-health disparities in countries such as Mexico. Further research is needed to monitor the longer-term consequences of the pandemic and school closures for youth. Population-level policies designed to provide economic protection and allow youth to return to school would have a favorable impact on the mental health and suicidal behavior of adolescents.

Statements

Statements of ethics

The study was conducted according to the Helsinki Declaration, and was approved by the Ethics and Research Committees of the National Institute of Public Health in Mexico.

Role of the funding sources

The authors received no specific funding for this work.

CRedit authorship contribution statement

Rosario Valdez-Santiago: Conceptualization, Writing – original draft, Writing – review & editing. **Aremis Villalobos:** Conceptualization, Formal analysis, Methodology, Writing – original draft, Writing – review & editing. **Luz Arenas-Monreal:** Writing – original draft, Writing – review & editing. **Catalina González-Forteza:** Writing – original draft, Writing – review & editing. **Alicia Edith Hermsillo-de-la-Torre:** Writing – original draft, Writing – review & editing. **Corina Benjet:** Writing – original draft, Writing – review & editing. **Fernando A. Wagner:** Methodology, Writing – original draft, Writing – review & editing.

Declaration of Competing Interest

The authors have no conflicts of interest regarding this research study.

Acknowledgement

None.

References

- Dubé, J.P., Smith, M.M., Sherry, S.B., Hewitt, P.L., Stewart, S.H., 2021. Suicide behaviors during the COVID-19 pandemic: a meta-analysis of 54 studies. *Psychiatry Res.* 301, 113998 <https://doi.org/10.1016/j.psychres.2021.113998>. Jul.
- Gutiérrez, J.P., 2013. Clasificación socioeconómica de los hogares en la ENSANUT 2012. *Salud Pública Mex.* 55, S341–S346.
- Gutiérrez-García, R.A., Benjet, C., Borges, G., Méndez Ríos, E., Medina-Mora, M.E., 2017. NEET adolescents grown up: eight-year longitudinal follow-up of education, employment and mental health from adolescence to early adulthood in Mexico city. *Eur. Child Adolesc. Psychiatry* 26 (12), 1459–1469. <https://doi.org/10.1007/s00787-017-1004-0>.
- Hannan, S., Honjo, K., Raissi, M., <https://www.imf.org/en/Publications/WP/Issues/2020/10/13/Mexico-Needs-a-Fiscal-Twist-Response-to-Covid-19-and-Beyond-49817>.
- Haw, C., Hawton, K., Gunnell, D., Platt, S., 2015. Economic recession and suicidal behaviour: possible mechanisms and ameliorating factors. *Int. J. Soc. Psychiatry* 61 (1), 73–81, 36.
- Hermosillo-de la Torre, A.E., González-Forteza, C., Rivera-Heredia, M.E., Méndez-Sánchez, C., González-Betanzos, F., Palacios-Salas, P., Jiménez, A., Wagner, F.A., 2020. Understanding suicidal behavior and its prevention among youth and young adults in Mexico. *Prev Med* 138, 1–6. ISSN e: 00917435.
- Hermosillo-de la Torre, A.E., Arteaga-de Luna, S.M., Acevedo-Rojas, L.D., Juárez-Loya, A., Jiménez-Tapia, J.A., Pedroza-Cabrera, J., González-Forteza, C., Cano, M., Wagner, F.A., 2021. Psychosocial correlates of suicidal behavior among adolescents under confinement due to the COVID-19 pandemic in Aguascalientes, Mexico. *Int. J. Environ. Res. Public Health* 18 (9), 4977. <https://doi.org/10.3390/ijerph18094977>.
- Instituto Nacional de Estadística Geografía e Informática. Encuesta Nacional de Ocupación y Empleo. 2021 <https://www.inegi.org.mx/contenidos/saladeprensa/aproposito/2020/trabajoNal.pdf>.
- Levinsohn, P.M., Rohde, P., Seeley, J.R., Baldwin, C.L., 2001. Gender differences in suicide attempts from adolescence to young adulthood. *J. Am. Acad. Child Adolesc. Psychiatry* 40 (4), 427–434. <https://doi.org/10.1097/00004583-200104000-00011>. AprPMID: 11314568.
- Nock, M.K., Green, J.G., Hwang, I., McLaughlin, K.A., Sampson, N.A., Zaslavsky, A.M., Kessler, R.C., 2013. Prevalence, correlates, and treatment of lifetime suicidal behavior among adolescents: results from the national comorbidity survey replication adolescent supplement. *JAMA Psychiatry* 70 (3). <https://doi.org/10.1001/2013.jamapsychiatry.55>. Mar300-10PMID: 23303463; PMCID: PMC3886236.
- Nordt, C., Warnke, I., Seifritz, E., Kawohl, W., 2015. Modelling suicide and unemployment: a longitudinal analysis covering 63 countries, 2000–11. *Lancet Psychiatry* 2 (3), 239–245.
- Pirkis, J., John, A., Shin, S., Delgado-Banos, M., Arya, V., Aguilar, P.A., Spittal, M.J., 2021. Suicide trends in the early months of the COVID-19 pandemic: interrupted time series analysis of preliminary data from 21 countries. *Lancet Psychiatry* 1–10. [https://doi.org/10.1016/S2215-0366\(21\)00091-2](https://doi.org/10.1016/S2215-0366(21)00091-2).
- Remus, R., 2020. Suicide in adolescence: A review of Literature. *Rev. Asistenta Soc.* 3, 109–120.
- Rivera-Rivera, L., Fonseca-Pedrero, E., Séris-Martínez, M., 2020. Prevalencia y factores psicológicos asociados con conducta suicida en adolescentes ENSANUT 2018-19. *Salud Pública Mex.* 62 (6), 672–681.
- Rockstroh, F., Reichl, C., Lerch, S., Fischer-Waldschmidt, G., Ghinea, D., Koenig, J., Resch, F., Kaess, M., 2021. Self-rated risk as a predictor of suicide attempts among high-risk adolescents. *J. Affect. Disord.* 282, 852–857. <https://doi.org/10.1016/j.jad.2020.12.110>. ISSN 0165-0327.
- Romero-Martínez, M., Shamah-Levy, T., Vielma-Orozco, E., Heredia-Hernández, O., Mojica-Cuevas, J., Cuevas-Nasu, L., Rivera-Dommarco, J., 2019. Encuesta nacional de salud y nutrición 2018–2019: metodología y perspectivas. *Salud Pública Mex.* 61, 917–923.
- Romero-Martínez, M., Barrientos-Gutiérrez, T., Cuevas-Nasu, L., Bautista-Arredondo, S., Colchero, M.A., Gaona-Pineda, E.B., Lazcano-Ponce, E., Martínez-Barnetteche, J., Alpuche-Aranda, C., Rivera-Dommarco, J., Shamah-Levy, T., 2021. Metodología de la encuesta nacional de salud y nutrición 2020 sobre Covid-19. *Salud Pública Mex.* 63, 444–451. <https://doi.org/10.21149/12580>.
- Valdez-Santiago, R., Villalobos, A., Arenas-Monreal, L., González-Forteza, C., Hermsillo-de la Torre, A.E., Benjet, C., Wagner, F.A., 2021. Comparative analysis of lifetime suicide attempts among Mexican adolescents over the past 12 years. *J. Environ. Public Health*. *Int. J. Environ. Res. Public Health* 2021, 18, 5419. <https://doi.org/10.3390/ijerph18105419>. ISSN 1660-4601.
- Xiong, J., Lipsitz, O., Nasri, F., Lui, L.M.W., Gill, H., Phan, L., Chen-Li, D., Iacobucci, M., Ho, R., Majeed, A., McIntyre, R.S., 2020. Impact of COVID-19 Pandemic on Mental Health in the General Population: A Systematic Review. *J. Affect. Disord.* 277, 55–64. <https://doi.org/10.1016/j.jad.2020.08.001>.
- World Health Organization, 2020. Coronavirus disease2019 (Covid-19) Situation Report-51. (accessed on 14 May 2021).